

NEW YORK, Sept. 11, 2013 /PRNewswire/ -- Reportlinker.com announces that a new market research report is available in its catalogue:

[EpiCast Report: Overweight and Obesity - Epidemiology Forecast to 2022](http://www.reportlinker.com/p01623033/EpiCast-Report-Overweight-and-Obesity---Epidemiology-Forecast-to-2022.html#utm_source=prnewswire&utm_medium=pr&utm_campaign=Pathology) http://www.reportlinker.com/p01623033/EpiCast-Report-Overweight-and-Obesity---Epidemiology-Forecast-to-2022.html#utm_source=prnewswire&utm_medium=pr&utm_campaign=Pathology

EpiCast Report: Overweight and Obesity - Epidemiology Forecast to 2022

Summary

Overweight and obesity are characterized by abnormal or excessive fat accumulation in the body, which can increase the likelihood of developing type 2 diabetes and cardiovascular diseases. Obesity is an escalating public health problem with an increasing worldwide prevalence and has become a well-known threat to public health. As obesity is a complex condition associated with a wide range of lifestyle, genetic, social, cultural, and environmental factors, its prevalence may vary across countries, depending on the variations in these contributing factors.

GlobalData's epidemiological analysis forecasts that prevalent cases of overweight in the 9MM (US, France, Germany, Italy, Spain, UK, Japan, Brazil, and Canada) will increase from 250.05 million in 2012 to 271.86 million cases in 2022 at an annual growth rate (AGR) of 0.86%. Approximately 57% of the 250.05 million prevalent cases of overweight adults in the 9MM in 2012 were in men. GlobalData epidemiologists forecast that there were 167.39 million prevalent

cases of obesity in the 9MM in 2012, which will increase to 213.34 million cases in 2022, increasing at an AGR of 2.75%. Approximately 52% of the 167.39 million prevalent cases of obese adults in the 9MM in 2012 were in women.

According to GlobalData's forecast, the prevalent cases of both overweight and obesity will increase dramatically over the next decade, with the number of obese adults increasing by an alarming 27.50%. Overweight and obesity have reached pandemic proportions; however, GlobalData epidemiologists believe that with the alarming increase in the prevalence of both overweight and obesity, the morbidity and mortality associated with the disease will continue to increase dramatically as well.

Scope

- The Overweight and Obesity EpiCast Report provides an overview of the risk factors, comorbidities, and global trends of overweight and obesity in the 9MM (US, France, Germany, Italy

, Spain

, UK,

Japan

, Brazil

, and

Canada

). It includes a 10-year epidemiology forecast of the prevalent cases of overweight, obesity, obesity class I, obesity class II, and obesity class III segmented by sex and age. In addition, the report includes a 10-year forecast of the prevalent cases of obesity-associated comorbidities, including diagnosed diabetes, diagnosed hypertension, and dyslipidemia, among adults with overweight/obesity in some of these markets.- The overweight and obesity epidemiology report is written and developed by Masters- and PhD-level epidemiologists.- The EpiCast Report is in-depth, high quality, transparent and market-driven, providing expert analysis of disease trends in the 9MM.

Reasons to buy

- Develop business strategies by understanding the trends shaping and driving the global

overweight and obesity market. - Quantify patient populations in the global overweight and obesity market to improve product design, pricing, and launch plans. - Organize sales and marketing efforts by identifying the age groups and sex that present the best opportunities for overweight and obesity therapeutics in each of the markets covered. - Identify the number of obesity class I, obesity class II, and obesity class III prevalent cases.

1 Table of Contents
1 Table of Contents
41.1 List of Tables
71.2 List of Figures
82 Introduction
102.1 Catalyst
102.2 Related Reports
113 Epidemiology
123.1 Disease Overview
123.2 Risk Factors and Comorbidities
133.2.1 Family history is a strong predictor of obesity
133.2.2 Physical inactivity is an independent predictor of obesity
143.2.3 Excessive caloric intake doubles the risk for obesity
153.2.4 Hypertension is as high as 42% in obese adults
163.2.5 Dyslipidemia and type 2 diabetes are common comorbidities in obese patients
173.3 Global Trends
193.3.1 US
203.3.2 EU
223.3.3 Japan
283.3.4 Brazil
293.3.5 Canada
313.4 Forecast Methodology
323.4.1 Sources Used
393.4.2 Forecast Assumptions and Methods: Prevalent Cases of Overweight, Obesity, Obesity by Class, and Comorbidities
463.4.3 Sources Not Used
633.5 Epidemiology Forecast of Overweight (2012–2022)
633.5.1 Prevalent Cases of Overweight
633.5.2 Age-Specific Prevalent Cases of Overweight
653.5.3 Sex-Specific Prevalent Cases of Overweight
673.5.4 Age-Standardized Prevalence of Overweight
703.6 Epidemiology Forecast of Obesity (2012–2022)
713.6.1 Prevalent Cases of Obesity
713.6.2 Age-Specific Prevalent Cases of Obesity
733.6.3 Sex-Specific Prevalent Cases of Obesity
753.6.4 Age-Standardized Prevalence of Obesity
763.7 Epidemiology Forecast of Obesity Class I (2012–2022)
773.7.1 Prevalent Cases of Obesity Class I
773.7.2 Age-Specific Prevalent Cases of Obesity Class I
793.7.3 Sex-Specific Prevalent Cases of Obesity Class I
813.7.4 Age-Standardized Prevalence of Obesity Class I
833.8 Epidemiology Forecast of Obesity Class II (2012–2022)
853.8.1 Prevalent Cases of Obesity Class II
853.8.2 Age-Specific Prevalent Cases of Obesity Class II
873.8.3 Sex-Specific Prevalent Cases of Obesity Class II
893.8.4 Age-Standardized Prevalence of Obesity Class II
913.9 Epidemiology Forecast of Obesity Class III (2012–2022)
933.9.1 Prevalent Cases of Obesity Class III
933.9.2 Age-Specific Prevalent Cases of Obesity Class III
953.9.3 Sex-Specific Prevalent Cases of Obesity Class III
973.9.4 Age-Standardized Prevalence of Obesity Class III
993.10 Epidemiology Forecast of Comorbidities among Adults with Overweight/Obesity (2012 and 2022)
1013.10.1 Prevalent Cases of Diagnosed Diabetes among Adults with Overweight/Obesity
1013.10.2 Prevalent Cases of Diagnosed Hypertension in Adults with Overweight/Obesity
1043.10.3 Prevalent Cases of Dyslipidemia among Adults with Overweight/Obesity
1063.11 Discussion
1093.11.1 Conclusions on Epidemiological Trends
1093.11.2 Limitations of the Analysis
1103.11.3 Strengths of the Analysis
1114 Appendix
1124.1 Bibliography
1124.2 About the Authors
1184.2.1 Epidemiologists
1184.2.2 Reviewers
1184.2.3 Global Director of Epidemiology and Health Policy
1194.2.4 Global Head of Healthcare
1204.3 About GlobalData
1214.4 About EpiCast
1214.5 Disclaimer
121

1.1 List of Tables

Table 1: The WHO Classification System of Adult Overweight and Obesity According to BMI
12
Table 2: Risk Factors and Comorbidities of Obesity
13
Table 3: 9MM, Age-Adjusted and Crude Prevalence (%) of Obesity, by Sex, Ages ≥20 Years, 2008
20
Table 4: JASSO and WHO Classifications of Obesity
28
Table 5: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Overweight
33
Table 6: Sources of Epidemiological Data Used for

Forecasting the Prevalent Cases of Obesity 34
Table 7: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Obesity Class I, Class II, and Class III 36
Table 8: Sources of Epidemiological Data Used for Forecasting the Prevalent Cases of Comorbidities in Overweight/Obese 38
Table 9: 9MM, Prevalent Cases of Overweight, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 64
Table 10: 9MM, Prevalent Cases of Overweight, By Age, Both Sexes, N (Millions), Row (%), 2012 66
Table 11: 9MM, Prevalent Cases of Overweight, Ages ?18 Years, By Sex, N (Millions), Row (%), 2012 68
Table 12: 9MM, Prevalent Cases of Obesity, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 72
Table 13: 9MM, Prevalent Cases of Obesity, By Age, Both Sexes, N (Millions), Row (%), 2012 74
Table 14: 9MM, Prevalent Cases of Obesity, Ages ?18 Years, by Sex, N (Millions), Row (%), 2012 75
Table 15: 9MM, Prevalent Cases of Obesity Class I, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 78
Table 16: 9MM, Prevalent Cases of Obesity Class I, by Age, Both Sexes, N (Millions), Row (%), 2012 80
Table 17: 9MM, Prevalent Cases of Obesity Class I, Ages ?18 Years, By Sex, N (Millions), Row (%), 2012 82
Table 18: 9MM, Prevalent Cases of Obesity Class II, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 86
Table 19: 9MM, Prevalent Cases of Obesity Class II, By Age, Both Sexes, N (Millions), Row (%), 2012 88
Table 20: 9MM, Prevalent Cases of Obesity Class II, Ages ?18 Years, By Sex, N (Millions), Row (%), 2012 90
Table 21: 9MM, Prevalent Cases of Obesity Class III, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 94
Table 22: 9MM, Prevalent Cases of Obesity Class III, By Age, Both Sexes, N (Millions), Row (%), 2012 96
Table 23: 9MM, Prevalent Cases of Obesity Class III, Ages ?18 Years, By Sex, N (Millions), Row (%), 2012 98
Table 24: 5MM*, Prevalent Cases of Diagnosed Diabetes among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 and 2022 102
Table 25: 6MM*, Prevalent Cases of Diagnosed Hypertension among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 and 2022 105
Table 26: 4MM, Prevalent Cases of Dyslipidemia* among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 and 2022 107
1.2 List of Figures

Figure 1: US, Overweight and Obesity Age-Adjusted Prevalence (%), Ages 20–74 Years, Men, 1960–2010 21
Figure 2: US, Overweight and Obesity Age-Adjusted Prevalence (%), Ages 20–74 Years, Women, 1960–2010 21
Figure 3: France, Overweight and Obesity Prevalence (%), Ages ?15 Years, Men, 1997–2006 23
Figure 4: France, Overweight and Obesity Prevalence (%), Ages ?15 Years, Women, 1997–2006 23
Figure 5: Italy, Overweight and Obesity Prevalence (%), Ages ?18, Men, 1983–2010 25
Figure 6: Italy, Overweight and Obesity Prevalence (%), Ages ?18, Men, 1983–2010 25
Figure 7: UK, Overweight and Obesity Prevalence (%), Ages ?16 Years, Men, 1993–2011 27
Figure 8: UK, Overweight and Obesity Prevalence (%), Ages ?16 Years, Women, 1993–2011 27
Figure 9: Japan, Obesity Prevalence (%), Ages ?20 Years, By Sex, 1976–2006 29
Figure 10: Brazil, Overweight and Obesity Prevalence (%), Ages ?20 Years, Men, 1975–2003 30
Figure 11: Brazil, Overweight and Obesity Prevalence (%), Ages ?20 Years, Men, 1975–2003 30

, Overweight and Obesity Prevalence (%), Ages ?20 Years, Women, 1975–2003 30Figure 12: Canada
, Obesity Prevalence (%), Ages ?18 Years, Both Sexes, 1978–2008 32Figure 13: 9MM, Prevalent Cases of Overweight, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 65Figure 14: 9MM, Prevalent Cases of Overweight, by Age, Both Sexes, N (Millions), 2012 67Figure 15: 9MM, Prevalent Cases of Overweight, Ages ?18 Years, by Sex, N (Millions), 2012 69Figure 16: 9MM, Overweight Age-Standardized Prevalence (%), Ages ?18 Years, By Sex, 2012 71Figure 17: 9MM, Prevalent Cases of Obesity, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 73Figure 18: 9MM, Prevalent Cases of Obesity, By Age, Both Sexes, N (Millions), 2012 74Figure 19: 9MM, Prevalent Cases of Obesity, Ages ?18 Years, by Sex, N (Millions), 2012 76Figure 20: 9MM, Obesity Age-Standardized Prevalence (%), Ages ?18 Years, By Sex, 2012 77Figure 21: 9MM*, Prevalent Cases of Obesity Class I, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 79Figure 22: 9MM*, Prevalent Cases of Obesity Class I, By Age, Both Sexes, N (Millions), 2012 81Figure 23: 9MM*, Prevalent Cases of Obesity Class I, Ages ?18 Years, by Sex, N (Millions), 2012 83Figure 24: 9MM*, Obesity Class I Age-Standardized Prevalence (%), Ages ?18 Years, by Sex, 2012 84Figure 25: 9MM*, Prevalent Cases of Obesity Class II, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 87Figure 26: 9MM*, Prevalent Cases of Obesity Class II, By Age, Both Sexes, N (Millions), 2012 89Figure 27: 9MM*, Prevalent Cases of Obesity Class II, Ages ?18 Years, by Sex, N (Millions), 2012 91Figure 28: 9MM*, Age-Standardized Prevalence (%) of Obesity Class II, Ages ?18 Years, By Sex, 2012 92Figure 29: 9MM*, Prevalent Cases of Obesity Class III, Ages ?18 Years, Both Sexes, N (Millions), 2012–2022 95Figure 30: 9MM*, Prevalent Cases of Obesity Class III, By Age, Both Sexes, N (Millions), 2012 97Figure 31: 9MM*, Prevalent Cases of Obesity Class III, Ages ?18 Years, By Sex, N (Millions), 2012 99Figure 32: 9MM*, Obesity Class III Age-Standardized Prevalence (%), Ages ?18 Years, By Sex, 2012 100Figure 33: 5MM*, Prevalent Cases of Diagnosed Diabetes among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 103Figure 34: 6MM*, Prevalent Cases of Diagnosed Hypertension among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 106Figure 35: 4MM*, Prevalent Cases of Dyslipidemia** among Adults with Overweight/Obesity, Both Sexes, N (Millions), 2012 108

To order this report: [EpiCast Report: Overweight and Obesity - Epidemiology Forecast to 2022](http://www.reportlinker.com/p01623033/EpiCast-Report-Overweight-and-Obesity---Epidemiology-Forecast-to-2022.html#utm_source=prnewswire&utm_medium=pr&utm_campaign=Pathology)
http://www.reportlinker.com/p01623033/EpiCast-Report-Overweight-and-Obesity---Epidemiology-Forecast-to-2022.html#utm_source=prnewswire&utm_medium=pr&utm_campaign=Pathology

Contact Clare: clare@reportlinker.com US: (339)-368-6001 Intl: +1 339-368-6001

SOURCE Reportlinker

EpiCast Report: Overweight and Obesity - Epidemiology Forecast to 2022

Written by Australian Business

RELATED LINKS <http://www.reportlinker.com>